

Differential effects of local habitat and landscape characteristics on bird communities in Mediterranean afforestations motivated by the EU Common Agrarian Policy

J. S. Sánchez-Oliver · J. M. Rey Benayas · L. M. Carrascal

Received: 6 May 2013 / Revised: 10 July 2013 / Accepted: 12 July 2013 / Published online: 2 August 2013
© Springer-Verlag Berlin Heidelberg 2013

Abstract We investigated the effects of local habitat structure and surrounding landscape characteristics (proportion of land use types and connectedness) on species density and composition of bird communities inhabiting the interior of young tree plantations on former cropland in central Spain, which were motivated by the Common Agrarian Policy. Variation of species density (number of species/0.78 ha) among tree plantations showed different environmental associations across seasons: local habitat was more important than landscape characteristics during winter, whereas they were similarly important during spring. Species density increased with the development of the tree layer in winter and with the presence of urban areas around tree plantations and cover of the herbaceous layer within them in the breeding season. We identified 15 species that exhibit high relative abundance in woodland habitats within the Mesomediterranean region of Central Spain that were absent in both seasons in the studied tree plantations, which were an attractive habitat for urban exploiter species but an unfavorable habitat for the regional forest species pool except for forest generalist species. Composition of bird assemblages was more related to local habitat structure

than to landscape characteristics around tree plantations and was rather similar in winter and spring seasons. The very different effects of local habitat and landscape characteristics on bird communities make difficult suggesting management practices with positive effects for all avifauna species during the entire year. We conclude that the small size and low maturity of the studied tree plantations do not contribute to enhancing the bird diversity value of current CAP aids to afforest former cropland with pines in the Mediterranean region.

Keywords Bird assemblages · Forest generalist species · Land use · Seasonal variation · Species density · Tree plantation

Introduction

Features of animal assemblages respond to the characteristics of both the local habitat and the landscape that surround such habitat, and these two sets of characteristics can interact affecting species composition and abundance (Fischer et al. 2011; Geiger et al. 2010; Moreno-Mateos et al. 2011; Piha et al. 2007; Wretenberg et al. 2010). Conversely, human activities may profoundly modify land cover and vegetation structure at both levels and, consequently, affect the composition and abundance of local communities (Blondel 1990; Heikkinen et al. 2004).

Large tracts of cropland and pastureland have been reforested in the world in recent decades by tree plantations or by secondary succession. Seven percent of the forest land is tree plantations at present, and their annual rate is growing as compared with afforestation by secondary succession (FAO 2011; Rey Benayas and Bullock 2012). These tree plantations have noticeable effects on both the abiotic environment and biological communities (Bremer and Farley 2010; Gómez-Aparicio et al. 2009; Munro et al. 2009; Poschlod et al. 2005), particularly on birds that are a taxonomic group of high

Communicated by C. Gortázar

Electronic supplementary material The online version of this article (doi:10.1007/s10344-013-0759-y) contains supplementary material, which is available to authorized users.

J. S. Sánchez-Oliver · J. M. Rey Benayas (✉)
Dpto. de Ciencias de la Vida, Edificio de Ciencias,
Universidad de Alcalá, 28871 Alcalá de Henares, Spain
e-mail: josem.rey@uah.es

J. S. Sánchez-Oliver
e-mail: juansalvador.sanchez@uah.es

L. M. Carrascal
Dpto. Biogeografía y Cambio Global, Museo Nacional
de Ciencias Naturales, MNCN-CSIC, José Gutiérrez Abascal, 2,
28006 Madrid, Spain
e-mail: mcnc152@mncn.csic.es

indicator value (Felton et al. 2010; Lindenmayer et al. 2010; Rey Benayas et al. 2010; Santos et al. 2006). In the European Union, the Common Agrarian Policy (CAP) has favored the transformation of farmland into tree plantations since 1993 by means of a scheme of aid for forestry measures in agriculture (EEC Council Regulation No. 2080/92), which has resulted on the afforestation of > 8 million ha to date (European Commission 2013a and 2013b). This afforestation program pursues both societal and environmental benefits, including control of erosion, prevention of desertification, regulation of the water regime, and increasing the fixation rate of carbon dioxide. The amount of afforested farmland will likely increase in a near future in some European regions because of subsidies to vineyard extirpation (e.g., 93,600 ha were extirpated in Spain in the 2008–2011 period of which 73.1 % belonged to La Mancha; Spanish Agrarian Guarantee 2012) together with subsidies to afforestation of former vineyards, which aim to ensure EU wine production matches demand and eliminate wasteful public intervention in EU wine markets (Regulation (EC) 479/2008).

Cropland afforestations in southern Europe are mostly based on coniferous species such as *Pinus halepensis* and *Pinus pinaster*. Afforested fields usually form an archipelago of man-made woodland habitat in the dominant agricultural matrix. These plantations may adversely affect open-habitat species that are of conservation concern in Europe, including birds, by replacing high-quality steppe habitat and increasing risk of predation (Cresswell 2008; Díaz et al. 1998; Reino et al. 2010). However, they may offer opportunities to woodland birds, providing suitable habitats for generalist species (Rey Benayas et al. 2010). Conversely, agricultural land abandonment and active afforestation should not be assumed to always benefit conservation, as it has been shown for birds of different biogeographic origin in agricultural lands of the Mediterranean region (increase in diversity with successional stage for Eurosiberian birds but not for Mediterranean species; Suárez-Seoane et al. 2002). Species–area relationships for bird communities in natural forests and pine plantations of Spain have been previously studied in detail (e.g., Díaz et al. 1998; Santos et al. 2006), demonstrating a very tight relationship between the area of forest islands and species richness. Nevertheless, little is known about how local species richness at standardized area units (i.e., species density) is affected by the surrounding landscape while taking into account habitat characteristics of the focal tree plantation.

In this study, we aim to assess the wintering and breeding bird communities in young tree plantations (<20 years old) motivated by the CAP that are embedded in Mediterranean agricultural landscapes of Central Spain. These plantations are located at the south-western limit of the Palaearctic, a region with impoverished woodland avifauna dominated by species of Mediterranean origin and woodland generalists (Carrascal and Díaz 2003; Monkkonen 1994; Tellería and Santos 1994),

and a strong seasonality in abiotic conditions and productivity that imposes widely different ecological scenarios throughout the year on the communities living in them (Newton 2007). They are established in small patches over a predominantly treeless landscape dominated by herbaceous or woody crops, where large mature forests of holm oaks that may serve as sources of woodland bird species are very scarce. Therefore, the avifauna in the plantations should be highly influenced by that inhabiting the surrounding landscape. This biogeographic scenario combined with the current CAP subsidies for afforestation on former arable land allow us testing the importance of local habitat characteristics and larger-scale features (e.g., the land cover surrounding the tree plantations) on bird assemblages. Moreover, the analysis of the responses of birds that colonize the interior of these afforestations in two contrasting seasons may proportionate insights about the temporal generality of their effects and suggest management practices that favor the implementation of friendly afforestation projects for woodland avifauna within deforested landscapes of the Mediterranean region on a seasonal basis.

Methods

Study area

Field work was conducted in tree plantations located in Campo de Montiel (La Mancha, situated in the southern Spanish plateau). The study area is ca. 440 km² within UTM coordinates x_1 , 4305423; x_2 , 4272951; y_1 , 458025; and y_2 , 483525 (zone 30S; Fig. 1 in Appendix 1). Altitude ranges between 690 and 793 m a.s.l. The climate is continental Mediterranean with dry and hot summers and cold winters. Mean annual temperature and total annual precipitation in the area during the last 30 years were 13.7 °C and 390 mm, respectively (retrieved from <http://www.aemet.es/>). These figures were 16.6 °C and 359.9 mm in 2011, when our bird surveys took place.

The area is a representative mosaic of different crops and semi-natural or introduced woody vegetation that is characteristic of large areas in Mediterranean landscapes. Croplands were mostly occupied by herbaceous crops (wheat and barley), harvested once a year in June, and permanent woody crops (olive trees, 3 to 5 m high and vineyards, 1 m high). Natural vegetation mostly consisted of holm oak *Quercus rotundifolia* L. woodland and riparian forests that have been mostly extirpated from this region. Until 1992, woodland cover was restricted to open holm oak patches, usually grazed by sheep and goats. However, as in many other Mediterranean landscapes, the agricultural land is subjected to intensive management (e.g., irrigation of vineyards and olive groves) and land use change. A major result of land use change is the abandonment of herbaceous cropland and vineyard extirpation and their

afforestation with the native Aleppo pine *P. halepensis* Mill. alone or mixed with holm oak and *Retama sphaerocarpa* (L.) Boiss., which has increased forest land in the last 20 years. These tree plantations are noticeably dominated by pines as they establish better and grow faster than the other planted species. Thus, height and diameter at breast height (dbh) of dominant pines are surrogates of the age of tree plantations.

Bird survey

First, all young forest plantations in the study area were identified using both orto-photos (Geographic Information System of Farming Land 2010; hereafter SigPac) and Google Earth®, and were later verified in the field. We found 99 plantations that were established in 1992 or later. Next, we selected the plantations to be surveyed for birds, excluding those smaller than 1 ha: 61 forest plantations with a mean area of 4.82 ha (SD=5.61; larger plantation=36.5 ha). Average spacing distance between studied plantations was 11.7 km (see Fig. 1 in Appendix 1). Pruning and thinning are the management practices performed on these plantations that modify their vegetation structure; 26 of our surveyed plantations were pruned and 16 of which were also thinned.

Species abundance and density were quantified by means of circular plot censuses that were carried out in winter (January and February) and spring (April and May) 2011, to study wintering and breeding bird communities, respectively. Every tree plantation was represented by one circular plot located at the center of the plantation. Census method consisted of point counts (Bibby et al. 2000), 10 min long each, recording all birds detected visually and/or acoustically within the 50-m radius plot (0.78 ha). We noted the presence of every bird species during the 10 min except if individuals were overflying the plot. Two censuses of each plot were carried out in each season, one in the morning between sunrise and 3 h later and one in the evening 2 h before sunset. The relative abundance of each species and local species density (i.e., number of species per 0.78 ha) were estimated using the average of the two censuses in each season. The small area covered by the plots, and the relatively long time devoted to bird counts (accumulated census time of 20 min in each season), maximizes the detection probability of species within the area of 0.78 ha and, thus, the accurate estimations of local species density and abundance (Shiu and Lee 2003). This time invested in bird census (25.6 min ha⁻¹) is considerably longer than that used in previous studies recording species richness in woodland islands (e.g., 10.2 min⁻¹ in pine plantations sampled by Díaz et al. 1998). Otherwise, our purpose was not to exhaustively characterize the avifauna of each plantation, but to analyze the variation of local species density in the interior of this novel habitat of an archipelago of young and small afforestations that punctuates the agricultural landscape. All

censuses were conducted by the same well-trained field ornithologist (JS S-O) on windless and rainless days.

To have a reference of the avifauna that potentially can colonize the studied plantations, we used the habitat breadth of the bird species in 15 main habitat categories as well as their relative abundance in woodlands within the Mesomediterranean region of Central Spain obtained from Carrascal and Palomino (2008).

Local habitat and landscape variables

We characterized two sets of variables related to tree plantations, namely (1) local habitat variables, which included vegetation in the bird census plots and area of the plantations and (2) landscape variables, which included tree plantation connectivity and land use around plantations.

1. Vegetation structure and composition of main plant species at each surveyed forest plantation were measured in 25- and 10-m radius plots that coincided with the centers of the bird census plots. This sampling was carried out before the spring bird census. In the 25-m radius plots, we directly counted or estimated by eye, after previous training, the following structural features of the vegetation: percentage cover of chamaephytes, shrubs and trees, average height of chamaephytes, shrubs and trees, and number of trunks <5, 5–10, 10–20, 20–40, and >40 cm in dbh. In the 10-m radius plots, we estimated percentage cover of herbs and bare soil and measured the average height of the herb layer. All vegetation measurements (Table 1) were carried out by the same observer (JS S-O) to avoid inter-personal bias.
2. Land use types were identified by means of land use layers taken from SigPac (see source above). They were analyzed with ArcGIS 10.0 in 1-km buffer-rings from the center of each forest plantation; on each buffer ring, the percentage of area occupied by each land use type was obtained, resulting in the figures shown in Table 1. Finally, for a target plantation, structural connectedness was measured as the average distance of the three closest plantations or natural woodland patches weighted by the area of such plantations or woodland patches (Table 1).

Statistical analyses

The effects of pruning on the development of the tree layer was tested by means of a MANOVA on percentage of tree cover, height of the tree layer, dbh, and number of trunks at >5 cm.

The relationships of species density and species composition with local habitat and landscape predictor variables were separately analyzed for the winter and the breeding season by means of Partial Least Squares Regressions (PLSR; Abdi

Table 1 Mean, standard deviation (SD), and range (min/max) of the local habitat and landscape variables describing the characteristics of the 61 studied tree plantations

	Mean	SD	Range	
Local habitat				
Area of tree plantation (ha)	4.8	5.6	1.0	36.5
Cover of the tree layer (%)	35.4	25.5	1.7	100
Average pine height (m)	3.5	1.5	0.9	7.2
Average trunk diameter of pines (cm dbh)	11.4	5.8	0	33.2
No. of pine trunks larger than 5 cm dbh/0.2 ha	70.5	50.7	0	185
Cover of the shrub layer (%)	4.7	8.8	0	46.2
Average height of the shrub layer (m)	1.2	1.1	0	3.3
Cover of the herbaceous layer (%)	54.3	40	0	100
Average height of the herbaceous layer (m)	0.4	0.3	0	1.1
Landscape around plantations				
Average distance to other woodlands (m)	739.7	621.7	14	2506
Streams, rivers, and lagoons (% cover)	0.7	1.1	0	4.1
Roads and rural tracks (% cover)	6.4	5.2	0	31.1
Woodlands (% cover)	4.2	4.7	0.1	25.2
Fruit groves (% cover)	1.1	1.3	0	5.4
Waste lands (% cover)	6.8	4.4	0	14.8
Olive groves (% cover)	21.9	23.7	0	94.7
Pastures with scattered trees (% cover)	0.4	1.6	0	9.4
Scrubland (% cover)	10.0	7.4	0	29.5
Pastures (% cover)	1.1	3.2	0	19.1
Dry herbaceous cropland (% cover)	18.2	9.2	0	40.8
Vineyards (% cover)	20.9	13.7	0	49.2
Vineyards with olive trees (% cover)	5.1	8.5	0	32.3
Dried fruit orchards (% cover)	0.6	2.4	0	16.9
Urban areas and scattered buildings (% cover)	2.4	4.2	0	25.8

2007). Sample units for these analyses were the 61 census plots in the tree plantations. Results obtained with PLSR are similar to those from conventional multiple regression techniques; however, PLSR allows for the simultaneous analysis of multiple response variables, and it is extremely robust to the effects of low sample size (i.e., overfitting) and high degree of correlation between predictor variables (i.e., severe multicollinearity) (Carrascal et al. 2009). PLSR establishes associations between the response variables and factors extracted from the predictor variables that maximize the explained variance in the response variables. These factors are defined as linear combinations of predictors, so the original multidimensionality is reduced to a lower number of orthogonal factors to detect structure in the relationships between predictor variables and between these factors and the response variables. The relative contribution of each predictor to the extracted factors was calculated by means of the square of predictor weights. The PLSR components regarding species

composition were obtained based on the abundance of those species with >0.1 birds/census plot; the abundance of 12 species in winter and 17 species in spring defined the response variables that were summarized in composition components by means of the linear combination of the species' abundances. Only those components significant after a tenfold validation procedure were retained (StatSoft 2011).

All statistical analyses were conducted in Statistica 10 (StatSoft 2011).

Results

Tree plantation and landscape characteristics

There was a broad variation in the local habitat variables of the studied tree plantations (Table 1). Overall, the number of pines at >5 cm dbh was not too large but there were a lot of small trees when considering the average trunk diameter of pines. Pruning enhanced the development of the tree layer according to a MANOVA (Wilk's $\lambda=0.752$, $p=0.003$, $n=61$).

There was also a considerable variation in the landscape characteristics around the tree plantations in an area mainly dominated by dry herbaceous cropland, olive tree groves, and vineyards (Table 1).

Species density

Average number of species per census plot of 0.78 ha did not significantly change between seasons (paired t test: $t=0.158$, $df=60$, $p=0.875$), being 4.38 species during winter time (range=0–9, SD=2.02, $n=61$ plots) and 4.43 species during the breeding season (range=1–10, SD=1.84). Winter and spring species density were not significantly correlated ($r=0.208$, $p=0.109$, $n=61$).

One significant component ($p<<0.001$) was obtained in each PLSR analysis of species density in the 61 studied tree plantations using all local habitat and landscape predictor variables, accounting for 31.9 and 31.4 % of total variance in winter and breeding season species density, respectively (Table 2; Fig. 1). Environmental effects on local species density were very different in both seasons. The weights of local habitat and landscape variables were not significantly correlated in winter and spring ($r=0.190$, $p=0.372$, $n=24$ predictor variables), thus defining different patterns of environmental determinism on species density in both seasons.

In winter, species density mainly increased with the development of the tree layer (cover, height and trunk diameter of pines), which was associated to low development of the herbaceous and shrub layers (Table 2; Fig. 1). None predictor variable describing landscape characteristics around the plantations attained a $|\text{weight}|>0.2$. Thus, the importance of local habitat variables on winter species density was considerably

Table 2 Results of the partial least squares regression (PLSR) models analyzing the variation in bird species density and bird species composition in 61 tree plantations during winter and the breeding season (spring) according to nine local habitat features of plantations and 15 landscape predictor variables

	SPP density		SPP composition	
	Winter	Spring	Winter	Spring
Local habitat				
Area of tree plantation (ha)	0.16	<i>-0.29</i>	0.02	0.01
Cover of the tree layer (%)	<i>0.39</i>	0.09	<i>0.42</i>	<i>0.41</i>
Average pine height (m)	<i>0.47</i>	0.12	<i>0.45</i>	<i>0.45</i>
Average trunk diameter of pines (cm dbh)	<i>0.41</i>	-0.03	<i>0.34</i>	<i>0.29</i>
No. of pine trunks larger than 5 cm dbh	<i>0.41</i>	-0.09	<i>0.41</i>	<i>0.33</i>
Cover of the shrub layer (%)	-0.06	<i>-0.39</i>	-0.16	-0.16
Average height of the shrub layer (m)	-0.22	<i>-0.34</i>	-0.28	-0.23
Cover of the herbaceous layer (%)	-0.10	0.19	-0.08	-0.21
Average height of the herbaceous layer (m)	-0.25	-0.03	-0.26	-0.20
Landscape around plantations				
Average distance to other woodlands (m)	-0.06	0.08	0.07	0.14
Streams, rivers, and lagoons (% cover)	0.08	-0.03	0.00	0.06
Roads and rural tracks (% cover)	0.17	-0.06	0.03	0.01
Woodlands (% cover)	-0.01	<i>-0.29</i>	-0.14	0.00
Fruit groves (% cover)	-0.15	<i>-0.34</i>	-0.12	-0.11
Waste lands (% cover)	0.01	<i>0.27</i>	-0.04	-0.17
Olive groves (% cover)	-0.07	0.08	-0.01	0.10
Pastures with scattered trees (% cover)	0.02	0.08	0.08	-0.04
Scrubland (% cover)	0.04	<i>-0.33</i>	-0.04	-0.04
Pastures (% cover)	0.01	0.02	0.08	-0.01
Dry herbaceous cropland (% cover)	-0.06	<i>-0.20</i>	-0.08	-0.11
Vineyards (% cover)	0.07	0.05	0.01	-0.04
Vineyards with olive trees (% cover)	0.06	0.19	0.19	<i>0.22</i>
Dried fruit orchards (% cover)	0.18	-0.13	<i>0.21</i>	0.09
Urban areas and scattered buildings (% cover)	-0.18	<i>0.26</i>	-0.16	-0.35

Figures shown are the predictor weights of each variable in each component (entries in italics are those with |weights|>0.2; this threshold was calculated according to the following equation: $(1/\text{no. of predictors})^{0.5}$)

higher than the importance of variables describing the landscape characteristics (calculated by means of the square of predictor weights), and was considerably higher than that expected considering the relative number of predictors in the two groups of variables (local habitat=0.86, landscape=0.14; the “null” proportions according to the number of predictors was 0.38 for nine local habitat variables and 0.62 for 15 landscape variables).




Fig. 1 Relationship between (a) the species density per 0.78 ha census plot of tree plantations in the winter (top) and (b) the breeding season (down) and the multivariate gradient (first PLSR component) defined by the PLSR analysis on nine local habitat and 15 landscape predictor variables

During the breeding season, species density was positively associated with the presence of waste lands, urban areas, and scattered buildings around them and negatively related to their size, the height, and cover of shrubs and the amount of area around plantations covered by woodland (mainly remaining patches of holm oak forests), fruit groves, scrubland, and dry herbaceous cropland (Table 2; Fig. 1). Characteristics of landscape surrounding the tree plantations were similarly important than local habitat in determining species density during the breeding season (summatory of the square of predictor weights: 0.42 for nine local habitat variables and 0.58 for 15 landscape characteristics, which were very similar to the “null” proportions of 0.38 and 0.62, respectively, according to the number of predictors).

Species composition

The avifauna was dominated by the great tit (*Parus major*), the chiffchaff (*Phylloscopus collybita*), the goldfinch (*Carduelis carduelis*), the wood pigeon (*Columba palumbus*), and the magpie (*Pica pica*) in wintertime and by the goldfinch, the spotless starling (*Sturnus unicolor*), the wood pigeon, and the magpie during the breeding season (spring; average of more than one detected individual per census plot in both seasons; Appendix 2).

The following species that exhibit high relative abundance in woodland habitats within the Mesomediterranean region of Central Spain according to Carrascal and Palomino (2008) were completely absent in both seasons in the studied tree plantations: great spotted woodpecker (*Dendrocopos major*), blackbird (*Turdus merula*), nuthatch (*Sitta europaea*), short-toed treecreeper (*Certhia brachydactyla*), firecrest (*Regulus ignicapillus*), coal tit (*Periparus ater*), crested tit (*Lophophanes cristatus*), long-tailed tit (*Aegithalos caudatus*), hawfinch (*Coccothraustes coccothraustes*), blue tit (*Cyanistes caeruleus*), rock bunting (*Emberiza cia*), jay (*Garrulus glandarius*), and Eurasian Hoopoe (*Upupa epops*). Similarly, other woodland species in the region such as robin (*Erithacus rubecula*) and Woodchat Shrike (*Lanius senator*) were very scarce in the studied plantations.

Relative abundances of species across the 61 studied tree plantations were not tightly correlated among themselves either in winter or during the breeding season, as defined by the low variance attained by the first components of the PLSRs in both seasons using the common species (those with more than 0.1 birds/plot): 7.9 % of variance in the relative abundances of 12 species in winter and 5.7 % of variance for 17 species in spring. Nevertheless, these loose patterns of co-variation in species abundances were highly associated with the plantation characteristics, mainly local habitat in both seasons (see below): $r=0.675$, $p<<0.001$ for winter and $r=0.700$, $p<<0.001$ for the breeding season.

The main pattern of co-variation in species abundances during the winter season was the association of the chiffchaff, great tit, magpie, wood pigeon, chaffinch (*Fringilla coelebs*), and goldfinch in tree plantations with a well developed tree layer (Fig. 2; see predictor variable weights in Table 2; these species were selected according to absolute values of loadings >0.2 in the component of species abundances). Conversely, there is a common pattern of increase in species abundances during the breeding season that associates the magpie, great tit, and wood pigeon in tree plantations with a tall and dense cover of the tree layer surrounded by relatively high cover of vineyard with olive trees, as opposed to the co-variation of abundances of rock pigeon (*Columba livia*), spotless starling, little bustard (*Tetrax tetrax*), and crested lark (*Galerida cristata*) in plantations with high cover of the shrubs and herb layers near urban areas (Fig. 2).

The importance of the environmental factors related to composition of bird assemblages was rather similar in winter and spring (Table 2), as the weights of local habitat and landscape variables were highly correlated in both seasons: $r=0.921$, $p<<0.001$, $n=24$ predictor variables). Moreover, the importance of local habitat variables in defining the co-variation of abundance of bird species was considerably higher than that of variables describing the landscape characteristics around tree plantations in both seasons calculated by means of the square of predictor weights (winter—local habitat=0.83, landscape=0.17; spring—local habitat=0.73, landscape=0.27; the “null” proportions according to the number of predictors were 0.38 and 0.62, respectively).


Discussion

Overall community composition

Our results show that the local composition of bird assemblages inhabiting the interior of young Mediterranean cropland afforestations are characterized by a few common dominant species, namely magpie, wood pigeon, and goldfinch in both seasons, great tit and chiffchaff in wintertime, and spotless starling in spring. These ubiquitous species are generalist birds of wooded areas, with broad geographical ranges and high population sizes in Spain (Carrascal and Palomino 2008; Martí and del Moral 2003). They are of little conservation concern in the European context (BirdLife International 2004). They are also of little sensibility to habitat fragmentation as they can thrive in very small woodland patches (Díaz et al. 1998; Razola and Rey Benayas 2009; Santos et al. 2002), such as those corresponding to the afforestations investigated in this study.

The biogeographical basis of the avifauna in this Mediterranean region, with an impoverished European forest avifauna dominated by species of early successional stages, probably limits the possibility of colonization of pure coniferous woodland species. Forest specialists of Mediterranean coniferous forests that require more mature and larger woodland patches (Díaz et al. 1998; Santos et al. 2006), such as the great spotted woodpecker, firecrest, crested tit, short-toed treecreeper, or nuthatch, were never recorded in these plantations, thus emphasizing the low suitability of these woodlands for forest avifauna of the region. This points to the importance of the biogeographic context when designing restoration plans with afforestations in agricultural-dominant landscapes (Suárez-Seoane et al. 2002) and enlightens the conflicts that can arise if single services of ecological restoration such as carbon sequestration by tree plantations are targeted without taking into account regional biodiversity (Bullock et al. 2011).

Fig. 2 Relationship between (a) the species composition of tree plantations in the winter (*top*) and (b) the breeding season (*down*) and the multivariate gradient (first PLSR component) defined by the PLSR analysis on nine local habitat and 15 landscape predictor variables



Relative effects of local habitat and landscape characteristics

The influence of different sets of environmental factors, namely local habitat of tree plantations and landscape characteristics, on bird communities changed considerably between seasons, with a prominent role of local habitat variables during winter for species density, and a more balanced importance of landscape characteristics around plantations and local habitat during the breeding season. During the breeding season, birds are spatially restricted to the focal place where they breed, and thus they show marked habitat preferences related to vegetation structure, which is an important attribute determining species composition of bird communities at the local scale (Hinsley et al. 2009; Hurlbert 2004). By contrast, during the winter period, birds

adopt a vagabonding lifestyle exploring a greater variety of habitats over larger areas to track the spatial and temporal distribution of food availability (Levey and Stiles 1992; Wiktander et al. 2001). From this perspective, local habitat should have a greater importance in the breeding season than in the winter in influencing bird communities of tree plantations within agricultural landscapes. Nevertheless, our results do not support this prediction for species density.

The negative influence of the area of tree plantations studied here on local species density is related to the fact that the probability of recording “ubiquitous/edge” bird species in the centre of plantations decreases as plantation area increases. This result, together with the remarkable negative influence of nearby woodlands on local species density in the interior of

the plantations, reinforces the idea of the low favorability of these young afforestations dominated by pines for the forest avifauna of the study region, especially if they are large.

The high importance of urban cover around the tree plantations on species density during spring points to the attractiveness of scarce woodland fragments to urban-exploiters of Central Spain (Palomino and Carrascal 2006), such as the rock dove (*C. livia*), collared dove (*Streptopelia decaocto*), greenfinch (*Carduelis chloris*), house sparrow (*Passer domesticus*), magpie, or spotless starling. It also emphasizes that urban development extends its impact on the surrounding habitats affecting bird communities, especially by the influence of just a few very common urban species (e.g., Findlay and Houlihan 1997; Odell and Knight 2001; Palomino and Carrascal 2007; Sauvajot et al. 1998). Urban and surrounding areas are a source of the ubiquitous and opportunistic nest predator magpie, and could thus entail additional conservation concern, because its overabundance around the cities could pose a deleterious effect on other bird species breeding in the plantations (e.g., Andren 1992; Groom 1993; Paradis et al. 2000). Similarly, Lindenmayer et al. (2012) found that another aggressive corvid reduced bird abundance in Australian tree plantations located in an agricultural landscape.

Management of tree plantations

The results of this study show that, overall, there are difficulties in making generalizations about the environmental factors that determine bird diversity inhabiting the interior of young tree plantations in Mediterranean agricultural landscapes on a year-round basis, and thus in outlining management recommendations to make them friendlier for the avifauna. These plantations offer opportunities for a few generalist forest bird species but are not perceived as an attractive breeding habitat for most forest species in the region. Furthermore, the youngest plantations with under-developed tree layer and presence of shrub and herbaceous layers benefit bird species that are characteristic of open farmland habitats such as the calandra lark, little bustard, and rock pigeon (Rey Benayas et al. 2010). As pruning of pines speeds up the development of the tree layer, a more generalized use of this practice would increase overall species density in winter and benefit forest species such as the wood pigeon, which is of interest to hunters, and insectivorous birds such as the great tit or blue tit, which have the potential of enhancing pest regulation in both tree plantations and crops around them (Jedlicka et al. 2011).

Conclusions

Local habitat and surrounding landscape characteristics in Mediterranean landscapes dominated by croplands had very different effects on bird communities inhabiting the interior of

young afforestations in the winter and breeding seasons, which make difficult suggesting extensive management practices with positive effects for all avifauna species during the entire year. These small, monotonous plantations are an attractive habitat for urban exploiter species but an unfavorable habitat for the regional forest species pool with the exception of the forest generalist species. Therefore, the small size and low maturity of the studied tree plantations do not contribute to enhancing the bird diversity value of current CAP aids to afforest former cropland with pines in the Mediterranean region. Further monitoring of bird communities as these plantations get older is necessary to provide more robust science-based management recommendations, and test the success of the implemented recommendation (more use of tree pruning) that the results of this study hinted.

Acknowledgments Projects from the Spanish Ministry of Science and Education (CGL2010-18312) and the Government of Madrid (S2009AMB-1783, REMEDINAL-2) are currently providing financial support for this body of research. Claire Jasinski kindly improved the English of an early draft. We are indebted to two anonymous reviewers whose comments improved a previous version of this manuscript.

References

- Abdi H (2007) Partial least square regression (PLS regression). In: Salkind NJ (ed) Encyclopedia of measurement and statistics. Sage, Thousand Oaks, pp 740–744
- Andren H (1992) Corvid density and nest predation in relation to forest fragmentation—a landscape perspective. *Ecology* 73:794–804
- Bibby C, Burgess ND, Hill DA, Mustoe SH (2000) Bird census techniques, 2nd edn. Academic, London
- BirdLife International (2004) Birds in Europe: population estimates, trends and conservation status. BirdLife International, Cambridge, UK
- Blondel J (1990) Biogeography and history of forest bird faunas in the Mediterranean zone. In: Keast A (ed) Biogeography and ecology of forest bird communities. SPB Academic Publishing, The Hague, pp 95–107
- Bremer LL, Farley KA (2010) Does plantation forestry restore biodiversity or create green deserts? A synthesis of the effects of land-use transitions on plant species richness. *Biodivers Conserv* 19:3893–3915
- Bullock JM, Aronson J, Newton AC et al (2011) Restoration of ecosystem services and biodiversity: conflicts and opportunities. *Trends Ecol Evol* 26:541–549. doi:10.1016/j.tree.2011.06.011
- Carrascal LM, Díaz L (2003) Asociación entre distribución continental y regional. Análisis con la avifauna forestal y de medios arbolados de la Península Ibérica. *Graellsia* 59:179–207
- Carrascal LM, Palomino D (2008) Las aves comunes reproductoras en España. Población en 2004–2006. SEO/BirdLife, Madrid, p 202
- Carrascal LM, Galván I, Gordo O (2009) Partial least squares regression as an alternative to current regression methods used in ecology. *Oikos* 118:681–690. doi:10.1111/j.1600-0706.2008.16881.x
- Cresswell W (2008) Non-lethal effects of predation in birds. *Ibis* 150:3–17
- Díaz M, Carbonell R, Santos T, Tellería JL (1998) Breeding bird communities in pine plantations of the Spanish plateau: biogeography, landscape and vegetation effects. *J Appl Ecol* 35:562–574
- European Commission (2013a) Forestry measures under the common agricultural policy. http://ec.europa.eu/agriculture/envir/report/en/forest_en/report.htm. Accessed 25 July 2013

- European Commission (2013b) EU agriculture - Statistical and economic information. http://ec.europa.eu/agriculture/statistics/agricultural/index_en.htm. Accessed 25 July 2013
- FAO (2011) State of the World's Forests 2011. Food and Agriculture Organization of the United Nations, Rome, p 179
- Felton A, Knight E, Wood J et al (2010) A meta-analysis of fauna and flora species richness and abundance in plantations and pasture lands. *Biol Conserv* 143:545–554. doi:10.1016/j.biocon.2009.11.030
- Findlay CS, Houlihan J (1997) Anthropogenic correlates of species richness in Southeastern Ontario wetlands. *Conserv Biol* 11:1000–1009. doi:10.1046/j.1523-1739.1997.96144.x
- Fischer C, Flohre A, Clement LW et al (2011) Mixed effects of landscape structure and farming practice on bird diversity. *Agric Ecosyst Environ* 141:119–125. doi:10.1016/j.agee.2011.02.021
- Geiger F, De Snoo GR, Berendse F et al (2010) Landscape composition influences farm management effects on farmland birds in winter: a pan-European approach. *Agric Ecosyst Environ* 139:571–577. doi:10.1016/j.agee.2010.09.018
- Geographic Information System of Farming Land (2010) Geographic information system of farming land. Available from <http://pagina.jccm.es/agricul/sigpac.htm>.
- Gómez-Aparicio L, Zavala MA, Bonet FJ, Zamora R (2009) Are pine plantations valid tools for restoring Mediterranean forests? An assessment along abiotic and biotic gradients. *Ecol Appl* 19:2124–2141. doi:10.1890/08-1656.1
- Groom DW (1993) Magpie *Pica pica* predation on Blackbird *Turdus merula* nests in urban areas. *Bird Study* 40:55–62. doi:10.1080/00063659309477129
- Heikkinen R, Luoto M, Virkkala R (2004) Effects of habitat cover, landscape structure and spatial variables on the abundance mosaic of birds in an agricultural-forest mosaic. *J Appl Ecol* 41:824–835
- Hinsley SA, Hill RA, Fuller RJ et al (2009) Bird species distributions across woodland canopy structure gradients. *Comm Ecol* 10:99–110. doi:10.1556/ComEc.10.2009.1.12
- Hurlbert AH (2004) Species-energy relationships and habitat complexity in bird communities. *Ecol Lett* 7:714–720. doi:10.1111/j.1461-0248.2004.00630.x
- Jedlicka JA, Greenberg R, Letourneau DK (2011) Avian conservation practices strengthen ecosystem services in California vineyards. *PLoS One* 6:e27347. doi:10.1371/journal.pone.0027347
- Levey DJ, Stiles FG (1992) Evolutionary precursors of long-distance migration: resource availability and movement patterns in Neotropical landbirds. *Am Nat* 140:447. doi:10.1086/285421
- Lindenmayer DB, Knight EJ, Crane MJ et al (2010) What makes an effective restoration planting for woodland birds? *Biol Conserv* 143:289–301. doi:10.1016/j.biocon.2009.10.010
- Lindenmayer DB, Northrop-Mackie AR, Montague-Drake R et al (2012) Not all kinds of revegetation are created equal: revegetation type influences bird assemblages in threatened Australian woodland ecosystems. *PLoS One* 7:e34527. doi:10.1371/journal.pone.0034527
- Martí R, Del Moral JC (2003) Atlas de las Aves Reproductoras de España. p. 733
- Monkkonen M (1994) Diversity patterns in Palearctic and Nearctic forest bird assemblages. *J Biogeogr* 21:183–195
- Moreno-Mateos D, Rey Benayas JM, Pérez-Camacho L et al (2011) Effects of land use on nocturnal birds in a Mediterranean agricultural landscape. *Acta Ornithologica* 46:173–182. doi:10.3161/000164511X625946
- Munro NT, Fischer J, Wood J, Lindenmayer DB (2009) Revegetation in agricultural areas: the development of structural complexity and floristic diversity. *Ecol Appl* 19:1197–1210. doi:10.1890/08-0939.1
- Newton I (2007) The migration ecology of birds. Academic, London
- Odell EA, Knight RL (2001) Songbird and medium-sized mammal communities associated with exurban development in Pitkin County, Colorado. *Conserv Biol* 15:1143–1150. doi:10.1046/j.1523-1739.2001.0150041143.x
- Palomino D, Carrascal LM (2006) Urban influence on birds at a regional scale: a case study with the avifauna of northern Madrid province. *Landsc Urban Plan* 77:276–290. doi:10.1016/j.landurbplan.2005.04.003
- Palomino D, Carrascal LM (2007) Threshold distances to nearby cities and roads influence the bird community of a mosaic landscape. *Biol Conserv* 140:100–109. doi:10.1016/j.biocon.2007.07.029
- Paradis E, Baillie SR, Sutherland WJ et al (2000) Large-scale spatial variation in the breeding performance of song thrushes *Turdus philomelos* and blackbirds *T. merula* in Britain. *J Appl Ecol* 37:73–87. doi:10.1046/j.1365-2664.2000.00547.x
- Piha M, Tiainen J, Holopainen J, Vepsäläinen V (2007) Effects of land-use and landscape characteristics on avian diversity and abundance in a boreal agricultural landscape with organic and conventional farms. *Biol Conserv* 140:50–61. doi:10.1016/j.biocon.2007.07.021
- Poschold P, Bakker JP, Kahmen S (2005) Changing land use and its impact on biodiversity. *Basic Appl Ecol* 6:93–98. doi:10.1016/j.baae.2004.12.001
- Razola I, Rey Benayas JM (2009) Effects of woodland islets introduced in a Mediterranean agricultural landscape on local bird communities. *Web Ecol* 9:44–53
- Reino L, Porto M, Morgado R et al (2010) Does afforestation increase bird nest predation risk in surrounding farmland? *Forest Ecol Manage* 260:1359–1366. doi:10.1016/j.foreco.2010.07.032
- Rey Benayas JM, Bullock JM (2012) Restoration of biodiversity and ecosystem services on agricultural land. *Ecosystems* 15:883–889. doi:10.1007/s10021-012-9552-0
- Rey Benayas JM, Galván I, Carrascal LM (2010) Differential effects of vegetation restoration in Mediterranean abandoned cropland by secondary succession and pine plantations on bird assemblages. *Forest Ecol Manage* 260:87–95. doi:10.1016/j.foreco.2010.04.004
- Santos T, Tellería JL, Carbonell R (2002) Bird conservation in fragmented Mediterranean forests of Spain: effects of geographical location, habitat and landscape degradation. *Biol Conserv* 105:113–125. doi:10.1016/S0006-3207(01)00210-5
- Santos T, Tellería JL, Diaz M, Carbonell R (2006) Evaluating the benefits of CAP reforms: can afforestations restore bird diversity in Mediterranean Spain? *Basic Appl Ecol* 7:483–495. doi:10.1016/j.baae.2005.11.001
- Sauvajot RM, Buechner M, Kamradt DA, Schonewald CM (1998) Patterns of human disturbance and response by small mammals and birds in chaparral near urban development. *Urban Ecosyst* 2:279–297
- Shiu HJ, Lee P (2003) Assessing avian point-count duration and sample size using species. *Zool Stud* 42:357–367
- Spanish Agrarian Guarantee Fund (2012) Spanish Agrarian Guarantee Fund. www.fega.es. Accessed 16 Apr 2012
- StatSoft (2011) Statistica 10 (data analysis software system).
- Suárez-Seoane S, Osborne PEP, Baudry J (2002) Responses of birds of different biogeographic origins and habitat requirements to agricultural land abandonment in northern Spain. *Biol Conserv* 105:333–344. doi:10.1016/S0006-3207(01)00213-0
- Tellería JL, Santos T (1994) Factors involved in the distribution of forest birds in the Iberian Peninsula. *Bird Study* 41:161–169. doi:10.1080/00063659409477216
- Wiktander U, Olsson O, Nilsson SG (2001) Seasonal variation in home-range size, and habitat area requirement of the lesser spotted woodpecker (*Dendrocopos minor*) in southern Sweden. *Biol Conserv* 100:387–395. doi:10.1016/S0006-3207(01)00045-3
- Wretenberg J, Pärt T, Berg Å (2010) Changes in local species richness of farmland birds in relation to land-use changes and landscape structure. *Biol Conserv* 143:375–381. doi:10.1016/j.biocon.2009.11.001